



HOUSTON LAKE MINING INC.

Management Discussion & Analysis

For the Year Ended March, 31, 2013

This Management Discussion & Analysis ("MD&A"), for Houston Lake Mining Inc. ("HLM" or the "Company"), is prepared with an effective date of March 31, 2013, unless otherwise indicated and should be viewed in conjunction with the Company's financial statements. Other continuous disclosure documents, including the Company's press releases and other quarterly and annual reports are available through its filings with the securities regulatory authorities in Canada at www.sedar.com ("SEDAR") and are also available on the Company's website www.houstonlakemining.com.

TABLE OF CONTENTS



1. GLOSSARY OF TECHNICAL INFORMATION	4
2. FORWARD-LOOKING/SAFE HARBOUR STATEMENT AND FAIR DISCLOSURE STATEMENT	8
3. HISTORICAL RESOURCE ESTIMATES	9
4. INTRODUCTION	9
5. ASSET OVERVIEW.....	9
6. PAK RARE METALS PROJECT	10
6.1 Project Description and Location	11
6.2 Accessibility, Climate, Local Resources, Infrastructure and Physiography	12
6.3 History.....	14
6.4 Geological Setting.....	14
i. Regional Geology.....	14
ii. Property Geology	14
iii. Pakeagama Lake Pegmatite Geology.....	15
6.5 Geological Target: Rare Metal Target (Tanco Mine)	16
6.6 Exploration Programs	16
i. Past Exploration	16
ii. Recent Exploration	16
7. WEST CEDARTREE GOLD PROJECT (WCGP).....	17
7.1 Sale of WCGP	17
8. TIB LAKE PGE PROJECT.....	18
8.1 Definitive agreement on option to purchase Tib Lake PGE Project with North American Palladium	18
8.2 Tib Lake PGM Property Details	18
9. EXPLORATION STANDARDS.....	19
10. EXPLORATION ACTIVITIES AFFECTING THE BALANCE SHEET	19
10.1 Results of Operations for the three-month period ended March 31, 2013.....	19
10.2 Results of operations for the year ended March 31, 2013.....	20
11. FINANCING ACTIVITIES	20
12. CAPITAL STOCK	20
13. LIQUIDITY AND WORKING CAPITAL	20
14. OFF-BALANCE SHEET ARRANGEMENTS	20
15. ASSET-BACKED COMMERCIAL PAPER	20
16. RELATED PARTY TRANSACTIONS	20
17. OUTLOOK	21
18. RISKS AND UNCERTAINTIES	21
i. Credit Risk.....	21
ii. Liquidity Risk.....	21
iii. Asset Exploration Risk	21
iv. Commodity Price Risk	22
v. Interest Rate Risk.....	22
vi. Currency Risk.....	22



vii.	Sensitivity Analysis	22
viii.	Management Risk	22
19.	ADDITIONAL DISCLOSURE FOR VENTURE ISSUERS WITHOUT SIGNIFICANT REVENUE	22
20.	EVALUATION OF DISCLOSURE CONTROLS AND PROCEDURES.....	22

TABLE OF FIGURES

Figure I: Houston's Key Exploration Projects	10
Figure II: Project Location Map	11
Figure III: PAK Rare Metals Project Claim Locations	12
Figure IV: Project Regional Geology	13
Figure V: Channel Samples & Detailed Geology.....	17
Figure VI: Basic Geology of the Tib Lake and Lac Des Iles Intrusions.....	19



1. GLOSSARY OF TECHNICAL INFORMATION

The estimated mineral reserves and mineral resources discussed herein have been calculated in accordance with the Canadian Institute of Mining, Metallurgy and Petroleum (“CIM”) – Definitions Adopted by CIM Council on December 11, 2005 (the “CIM Standards”) which were adopted by the Canadian Securities Administrators’ National Instrument 43-101 Standards of Disclosure for Mineral Projects (“NI 43-101”). The following definitions are reproduced from the CIM Standards:

The term “**mineral reserves**” means the economically mineable part of a measured or indicated mineral resource demonstrated by at least a preliminary feasibility study. This study must include adequate information on mining, processing, metallurgical, economic and other relevant factors that demonstrate, at the time of reporting, that economic extraction can be justified. A mineral reserve includes allowances for dilution and losses that may occur when the material is mined. A “**proven mineral reserve**” is the economically mineable part of a measured mineral resource demonstrated by at least a preliminary feasibility study. This study must include adequate information on mining, processing, metallurgical, economic and other relevant factors that demonstrate, at the time of reporting, that economic extraction is justified. A “**probable mineral reserve**” is the economically mineable part of an indicated mineral resource, and in some circumstances a measured mineral resource, demonstrated by at least a preliminary feasibility study. This study must include adequate information on mining, processing, metallurgical, economic and other relevant factors that demonstrate, at the time of reporting, that economic extraction can be justified.

The term “**mineral resources**” means a concentration or occurrence of natural, solid, inorganic or fossilized organic material in or on the earth’s crust in such form and quantity and of such a grade or quality that it has reasonable prospects for economic extraction. The location, quantity, grade, geological characteristics and continuity of a mineral resource are known, estimated or interpreted from specific geological evidence and knowledge. A “**measured mineral resource**” is that part of a mineral resource for which quantity, grade or quality, densities, shape and physical characteristics are so well established that they can be estimated with confidence sufficient to allow the appropriate application of technical and economic parameters, to support production planning and evaluation of the economic viability of the deposit. The estimate is based on detailed and reliable exploration, sampling and testing information gathered through appropriate techniques from locations such as outcrops, trenches, pits, workings and drill holes that are spaced closely enough to confirm both geological and grade continuity. An “**indicated mineral resource**” is that part of a mineral resource for which quantity, grade or quality, densities, shape and physical characteristics can be estimated with a level of confidence sufficient to allow the appropriate application of technical and economic parameters, to support mine planning and evaluation of the economic viability of the deposit. The estimate is based on detailed and reliable exploration and test information gathered through appropriate techniques from location such as outcrops, trenches, pits, workings and drill holes that are spaced closely enough for geological and grade continuity to be reasonably assumed. An “**inferred mineral resource**” is that part of a mineral resource for which quantity and grade or quality can be estimated on the basis of geological evidence and limited sampling and reasonably assumed, but not verified. The estimate is based on limited information and sampling gathered through appropriate techniques from locations such as outcrops, trenches, pits, workings and drill holes.

The following technical terms may be used in this MD&A, and may appear capitalized or in lower case, without any difference in meaning:

Adamellite: Quartz monzonite. A medium to coarse-grained plutonic rock containing major plagioclase, orthoclase, and quartz, with minor biotite, hornblende, and accessory apatite, zircon, and opaque oxides.

Advance royalty - A form of royalty where the payment is made before the start of commercial production.

Albite – A plagioclase sodium feldspar.

Alkalic - Containing either sodium or potassium.

Alteration - Any change in the mineral composition of a rock that is brought about by physical or chemical means.

Amblygonite- A natural fluorophosphate of aluminum and lithium having the approximate formula, $2\text{LiF}\cdot\text{Al}_2\text{O}_3\cdot\text{P}_2\text{O}_3$. It should contain 10.1% Lithia, but actual samples average 8.2% due to partial replacement of the Lithia by soda and potash, partial alteration of the mineral to nonlithium minerals and the presence of impurities. It is the least expensive source of alumina-phosphate and is the highest Lithia containing lithium minerals. It produces opacity in glass dinnerware. Its usage is restricted due to its relatively limited availability.

Amphibolite grade metamorphism- A metamorphic facies assemblage of minerals formed at moderate to high pressures between 450 and 700°C during regional metamorphism.

Andesite – Igneous rock of intermediate composition.

Ankerite - An iron rich carbonate mineral.

Anomalous rare earth elements (Pakeagama)- Lithium (Li), Rubidium (Rb), Cesium, Beryllium (Be), Tantalum (Ta), columbium (Cb), Niobium (Nb), Tin (Sn), Gallium (Ga), Germanium (Ge), Hafnium (Hf).



Anomaly - Geochemical and/or geophysical data, which deviates from the norm.

Aplite- A light-coloured igneous rock characterized by a fine-grained saccharoidal (i.e., aplitic) texture. Aplites may range in composition from granitic to gabbroic, but the term aplite with no modifier is generally understood to mean granitic aplite, consisting essentially of quartz, potassium feldspar, and acid plagioclase. The term, from a Greek word meaning simple, was in use before 1823.

Archean - Oldest rocks of the Precambrian Era, older than about 2.5 billion years.

Argillaceous- Pertaining to, largely composed of, or containing clay-size particles or clay minerals, such as an argillaceous ore in which the gangue is mainly clay; esp. said of a sediment (such as marl) or a sedimentary rock (such as shale) containing an appreciable amount of clay.

Argillite- A compact rock, derived either from mudstone (claystone or siltstone), or shale, that has undergone a somewhat higher degree of induration than mudstone or shale but is less clearly laminated and without its fissility, and that lacks the cleavage distinctive of slate.

Assay - An analysis to determine the presence, absence or quantity of one or more chemical components.

Au - Chemical symbol for the element gold.

Basalt - Common dark and fine grained extrusive mafic volcanic rock.

Base Metal - A metal, such as copper, lead, nickel, zinc or cobalt.

Belt - A specific elongate area defined by unique geologic characteristics.

Beryl- A hexagonal mineral, $\text{Be}_3\text{Al}_2\text{Si}_6\text{O}_{18}$; green, blue-green, and other pale tints; in granite pegmatites, mica schists, and an accessory mineral in felsic igneous rocks; the chief source of beryllium. Transparent and coloured gem varieties include emerald, aquamarine, morganite, heliodor, golden beryl, bixbite, and vorobievite.

Breccia - Rock fragmented into angular components surrounded by a mass of finer grained material.

Bronzite - An orthorhombic mineral (pyroxene) consisting of a ferriferous variety of Enstatite and often having a lustre like that of bronze.

Carbonate - Mineral calcium carbonate (CaCO_3) and often a rock composed principally thereof.

Chalcopyrite - Copper iron sulphide (CuFeS_2).

Chlorite - A green platy iron-magnesium rich metamorphic mineral.

Claim (Mineral) - The area that confers mineral exploration/exploitation rights to the registered holder under the laws of the governing jurisdiction.

Clinopyroxene - Pyroxenes that crystallize in the monoclinic system and are commonly greenish in colour.

Collar - The top of a drill hole.

Columbite, Tantalite, Niobite- A natural oxide of niobium, tantalum, ferrous oxide, and manganese found in granites and pegmatite $[(\text{Fe},\text{Mn})(\text{Nb},\text{Ta})^2 \text{O}_6]$. Some tin or tungsten may be present in the mineral. Iron black to brownish black in colour; streak, dark red to black; luster, submetallic; Moh's hardness, 6; specific gravity, 5.2 to 7.9. See also: tantalite.

Columbium (CB)- Also called niobium (Nb). A shiny, white, soft, and ductile metallic element. Symbol, Nb (niobium) or Cb (columbium). The name niobium was adopted by the International Union of Pure and Applied Chemistry. Many chemical societies and government organizations refer to it as niobium, but most metallurgists, metals societies, and commercial producers still refer to the metal as columbium. Found in niobite (or columbite), niobite-tantalite, pyrochlore, and euxenite. Used as an alloying agent in carbon and alloy steels, in nonferrous metals, and in superconductive magnets.

Conglomerate - A sedimentary rock composed of rounded to subrounded transported fragments greater than 2 millimetres (pebbles, cobbles, boulders) cemented into a solid mass.

Dacitic - Igneous rock intermediate in compositions between andesite and rhyolite.

Diamond Drilling/Drill Hole - A method of obtaining a cylindrical core of rock by drilling with a diamond impregnated bit.

Diabase - A common basic igneous rock usually occurring in dykes or sills.

Diopside - A calcium-magnesium silicate (clinopyroxene), $\text{CaMg}(\text{Si}_2\text{O}_6)$, that is coloured white to light-green. The colour deepens with the addition of iron. Moh's hardness: 5 to 6; specific gravity: 3.2 to 3.3. It is found in regionally metamorphic rocks.

Dip - The angle at which a stratum is inclined from the horizontal.

Dyke - A tabular body of igneous rock cross cutting the host strata at a high angle.

Epithermal - A hydrothermal deposit formed close to surface at low temperature and pressure.

Elbaite- A trigonal mineral, $3[\text{Na}(\text{Li},\text{Al})_3\text{Al}_6(\text{OH},\text{F})_4(\text{BO}_3)_3\text{Si}_6\text{O}_{18}]$; tourmaline group; occurs in triangular and hexagonal prisms; varicolored; commonly zoned, pyroelectric and piezoelectric; in granites and granite pegmatites; and used as a gemstone (pink rubellite, blue indicolite, green verdolite, colourless achroite, zoned pink-white-green watermelon tourmaline).

Enstatite - A pyroxene mineral, $\text{MgO} \cdot \text{SiO}_2$. Its colour is yellowish- or greenish - grey.

Fault - A fracture in a rock along which there has been relative movement between the two sides either vertically or horizontally.

Fe - Chemical symbol for the element iron.

Feldspar - A group of common aluminosilicate minerals.

Felsic - Igneous rock composed principally of feldspars and quartz.

Ferriferous - A synonym for ferruginous (iron bearing).



Fertile - As in fertile granite. For example, a parental rock that gives rise to rare metal pegmatite.

Fluvial/fluviatile - Sedimentary material found in river beds.

Fold - Bend in strata or any planar structure.

Foliation - Parallel orientation of platy minerals or mineral banding in rocks.

Footwall - The wall or rock on the underside of a vein or structure.

Formation - A body of rock identified by lithological characteristics and stratigraphic position.

Fracture - A break in the rock, the opening of which allows the entry of mineral-bearing solutions.

Fuchsite - Mica with a characteristic (emerald) green colour arising from the presence of chrome or vanadium.

Gabbro - A fine to coarse grained, dark coloured crystalline igneous intrusive rock composed mainly of calcic plagioclase, clinopyroxene and sometimes olivine.

Gabbro-Norite - An igneous rock that is made up mainly of clinopyroxene and orthopyroxene and can contain upwards of 1/3 feldspar.

Geochemistry/Geochemical - Study of variation of chemical elements in rocks or soil.

Geology/Geological - Study of the Earth's history and life, mainly as recorded in rocks.

Geophysics/Geophysical - Study of the Earth by quantitative physical methods, either by surveys conducted on the ground, in the air (by fixed wing aircraft or helicopter) or in a borehole or drillhole.

Gold - A heavy, soft, ductile, malleable precious metal used in jewelry, dentistry, electronics and as an investment.

Grams per tonne (g/t) - A unit of measurement commonly used to quantify the concentration of precious metals.

Granoblastic: Pertaining to a homeoblastic type of texture in a nonschistose metamorphic rock upon which recrystallization formed essentially equidimensional crystals with normally well sutured boundaries.

Granophyre- An irregular microscopic intergrowth of quartz and alkali feldspar.

Granophyric- Of or pertaining to granophyres.

Greenstone belt - Area underlain by metamorphosed volcanic and sedimentary rocks, usually in a continental shield.

Greywacke - Grey sandstone consisting of poorly sorted grains of quartz, feldspar and rock fragments in a clay matrix.

Hangingwall - The wall or rock on the upper side of a vein or structure.

Hectare - A square of 100 metres on each side.

Hematite - Black to reddish brown, non-magnetic iron oxide (Fe_2O_3).

Holmquistite- An orthorhombic mineral, $\text{Li}_2(\text{Fe,Mg})_3\text{Al}_2\text{Si}_8\text{O}_{22}(\text{OH})_2$; amphibole group, with $\text{Mg}/(\text{Mg}+\text{Fe}^{2+}) = 0.1$ to 0.89; in granite pegmatites. A bluish-black silicate of lithium, magnesium, iron and aluminum with fluorine and hydroxyl. A lithium-bearing variety of glaucophane (or amphibole). Monoclinic, prismatic.

Holocrystalline - Rocks consisting entirely of crystalline minerals and no glass. The minerals may or may not have crystal boundaries, and the rocks may be granular or porphyritic.

Horizon - A defined layer within a stratigraphic sequence, having unique characteristics distinguishing it from the rest of the sequence.

Igneous - A classification of rocks formed from the solidification from a molten state.

Ilvaite- An orthorhombic and monoclinic mineral, $\text{CaFe}_2\text{FeSi}_2\text{O}_7\text{O}(\text{OH})$; in prisms with vertically striated faces; compact, massive, or fibrous; in some magnetite orebodies, in zinc and copper ores, in contact deposits in dolomitic limestone, and in sodalite syenite near Julianehaab, Greenland.

Infill drilling - Any method of drilling intervals between existing holes, used to provide greater geological detail and to help establish resource/reserve estimates.

Intrusive/Intrusions - An igneous rock that invades older rocks.

Iron formation (banded) - Chemically precipitated rock consisting of repeated thin layers of chert (silica) and iron oxides commonly magnetite and/or hematite.

Ironstone - A sedimentary rock containing a substantial proportion of iron.

IP/Induced polarization - Method of ground geophysical surveying employing an electrical current to determine indications of mineralization through the measurement of resistivity and chargeability.

JV/Joint venture - business arrangement usually between companies that defines each parties vested interest in an asset.

Komatiite - A volcanic rock containing a high concentration of magnesium and generally a low concentration of silica.

Lepidolite, lithia mica- A fluorosilicate of potassium, lithium and aluminum, $\text{K}^2\text{Li}^3\text{Al}^2(\text{AlSi}^3\text{O}^{10})^2(\text{O,OH,F})_4$; monoclinic; colour, pink, lilac and grayish-white; luster, pearly; perfect micaceous cleavage; comparatively rare mineral found in pegmatite dykes. Usually associated with other lithium-bearing minerals such as pink and green tourmaline, amblygonite and spodumene. Used as a source of lithium and in heat-resistant glass.

Mafic - An igneous rock composed chiefly of dark iron and manganese silicate minerals.

Magnetic Survey - A geophysical survey conducted on the Earth's surface that measures variations in the Earth's magnetic field caused by variations in rock type or geological structures.

Magnetite - Black, magnetic iron ore, an iron oxide (Fe_3O_4).

Mapping - The art and science of recording geological observations on a map.



Massive - Solid (without fractures) wide (thick) rock unit.

Melanocratic – Sixty to 100% of the (igneous) rock is made up of dark minerals.

Metamorphism/Metamorphic/Meta - A process whereby the composition of rock is modified by heat and pressure/A class of rock affected by metamorphism.

Mg - Chemical symbol for the element magnesium.

Mineralization - The concentration of metals and their chemical compounds in a body of rock.

Mineral Indicators for Rare Earths: blue fluorapatite, deep green elbite, high rubidium in feldspar, presence of pollucite, efficient zonal process, see anomalous elements.

Molybdenite - Molybdenum sulphide (MoS₂)

Mudstone – A fine grained sedimentary rock originally composed of clay and mud.

Mylonite- A hard, compact rock with a banded or streaky structure produced by extreme granulation in both strike-slip and thrust fault zones.

Niobium- see columbium.

NSR - Net Smelter Royalty – Royalty based on the actual gold sale price received less the cost of refining

Ore - Rock containing mineral(s) or metals that can be economically extracted to produce a profit.

Orogen/Orogeny – Deformation of a belt of rocks through folding and faulting, in many places accompanied by metamorphic and intrusive rocks that form mountains/the process of mountain building.

Orthopyroxene – Pyroxenes that crystallize in the orthorhombic system and are commonly brownish in colour.

Outcrop - An exposure of bedrock at the surface.

Pelite- A sediment or sedimentary rock composed of the finest detritus (clay- or mud-size particles); e.g., a mudstone, or a calcareous sediment composed of clay and minute particles of quartz. The term is equivalent to the Latin-derived term lutite.

Pelitic- Pertaining to or characteristic of pelite; especially said of a sedimentary rock composed of clay, such as a pelitic tuff representing a consolidated volcanic ash consisting of clay-size particles.

Peraluminous- Said of igneous rocks in which the molecular proportion of alumina exceeds that of soda, potash, and lime combined.

Peridotite – A general term for essentially non-feldspathic plutonic rocks consisting of olivine, with or without other mafic minerals. The other mafic minerals maybe amphiboles, pyroxenes or, less likely, micas. Minerals of the spinel group may be present.

Petalite- A monoclinic mineral, LiAlSi₄O₁₀; perfect cleavage; vitreous; resembles spodumene; a source of lithium salts; in granite pegmatites.

Pillowed - Volcanic rock texture that formed from the bulbous cooling of magma when cooled quickly in water.

Plunge - The vertical angle an ore body makes between the horizontal plane and the direction along which it extends, longitudinally to depth.

Pluton - Body of rock exposed after solidification at great depth.

Pollucite- An isometric mineral, (Cs, Na)₂Al₂Si₄O₁₂·H₂O; zeolite group; forms a series with analcime; colourless; occurs in granite pegmatites; a source of cesium and a minor gemstone.

ppb – Concentration in parts per billion.

ppm – Concentration in parts per million.

Porphyry - A rock consisting of larger crystals embedded in a more compact finer grained groundmass.

Prospecting – The art and science of searching for mineral deposits.

Proterozoic - The youngest part of the Precambrian from 2450 - 570 million years ago.

Pyrite - Iron sulphide mineral (FeS₂).

Pyroxene – A calcium/sodium ferromagnesium silicate.

Pyroxenite – A coarse-granited, holocrystalline rock consisting mainly of pyroxenes. Accessories may include biotite, hornblende and olivine.

Pyrrhotite – A slightly to moderately magnetic sulphide or iron that is often associated with the nickel sulphide, pentlandite. The mineral can contain nickel, and the elevated values for nickel in the gabbros of Tib Lake may be indicative of this. Nickel is a potential indicator element, along with copper, for Pt/Pd mineralization. The determination of distribution patterns for nickel and copper, especially, maybe a key to locating Pt/Pd mineralization.

Quartz - A mineral composed of silicon dioxide.

Rare Earth Elements; Rare Earths- A group of rare metals very similar to one another and to aluminum in many properties. Thorium is very closely associated in nature with this group, and the source of thorium salts is monazite.

Rare Earth Metals - A group of widely distributed metals comprising Scandium (At. No. 21), Yttrium (At. No. 39), and 15 elements of atomic numbers 57 to 71, inclusive. They have the same arrangement of the two external shells of electrons in their atoms and resemble one another very closely in chemical and physical properties, being thus most difficult to separate from each other. It is to this, rather than their actual rarity in nature, that they so described. The metals are divided into three groups (cerium, terbium and yttrium) based upon the methods used to separate them from their ores.

Rare Earths - A series of widely distributed but relatively scarce minerals, principally oxides of the rare earth metals.



Rhyolite – Igneous rock of felsic (silica rich) composition.

Rubellite- A pink gem variety of elbaite.

Sausseritization - The replacement of (plagioclase in particular) in basalts and gabbros by a fine-grained aggregate of zoisite, epidote, calcite, sericite and zeolite. The iron-magnesium minerals may be chloritized during the process.

Sandstone – A sedimentary rock composed mainly of sand-sized quartz and/or feldspar.

Schist – Rocks of medium-grade metamorphism with well developed lamellar minerals.

Sediment - Solid material that has settled down from a state of suspension in a liquid; may be transported and deposited by wind, water or ice, chemically precipitated from solution, or secreted by organisms, forms in layers in loose unconsolidated form.

Sedimentary - Pertaining to or containing sediment or formed by its deposition.

Sericite - Generally light coloured iron, magnesium and sodium rich mica.

Shear - A planar zone of deformed rock caused by the movement of the rock.

Siliceous - A rock rich in silica.

Sill - A tabular body of igneous rock conforming to the strata it invades.

Siltstone – A sedimentary rock with an intermediate grain size finer than sandstone with a higher clay fraction.

Soil Sampling - Systematic collection of soil samples from a series of different locations in order to study the distribution of its geochemical composition.

Specific gravity - The density of a substance relative to the density of water.

Splay – Branch of a fault.

Spodumene- A monoclinic mineral, $\text{LiAlSi}_2\text{O}_6$; pyroxene group; prismatic cleavage; in granite pegmatites in crystals up to scores of meters long (called logs); a source of lithium; may be of gem quality (lavender kunzite, green hiddenite). Formerly called triphane.

Stockwork – A local higher density of veins/stringers at numerous orientations

Strike - Direction or trend of a geologic structure.

Stringer - A very small vein or irregular filament of mineral(s) cutting a rock mass, occurs independently or as a branch of a larger vein.

Structure/Structural - Pertaining to geological structure such as folds, faults, etc.

Sulphide/Sulphidation - A group of minerals in which one or more metals are found in combination with sulphur/rock that has been sulphidized.

Syenite - An felsic intrusive igneous rock composed chiefly of the mineral orthoclase.

Tantalite- A mineral series ferrotantalite-manganotantalite; unless specified it refers to ferrotantalite, an orthorhombic mineral, FeTa_2O_6 ; black; in pegmatites; the main source of tantalum.

Tantalum (Ta): A rather brittle, lustrous, hard, heavy, grey metallic element. Symbol Ta. Occurs principally in the mineral columbite-tantalite, $(\text{Fe,Mn})(\text{Nb,Ta})_2\text{O}_6$. Widely used to fabricate chemical process equipment, nuclear reactors, and aircraft and missile parts. Used to make electrolytic capacitors, vacuum furnace parts, and surgical appliances.

Tonalite - A coarse-grained igneous rock, quartz-mica diorite. Two varieties are distinguished: soda tonalite, with albite in excess of anorthite, and lime tonalite, with anorthite in excess of albite. Compare to ademellite. They are biotitic at Pakeagama

Tholeiite – Mafic volcanic rock with higher silica and lower sodium, potassium and magnesium content.

Tourmaline - Any member of the trigonal mineral group, $\text{XY}_3\text{Z}_6(\text{BO}_3)_3\text{Si}_6\text{O}_{18}(\text{OH},\text{F})_4$ where X is Na partially replaced by Ca, K, Mg, or a vacancy, Y is Mg, Fe^{2+} , Li, or Al, and Z is Al and Fe^{3+} ; forms prisms of three, six, or nine sides; commonly vertically striated; vari-coloured; an accessory in granite pegmatites, felsic igneous rocks, and metamorphic rocks. Transparent and flawless crystals may be cut for gems.

Tuff/Pyroclastics - A rock formed of compacted volcanic fragments.

Turbidite - Submarine landslide along a continental slope containing large masses of sediment.

Ultramafic – A dark coloured igneous rock with a low silica content and characterized by mafic minerals, such as olivine, amphibole and pyroxene.

Unconformity - A surface of erosion that separates younger rocks from older rocks.

Uralitization – The process whereby primary pyroxene → uralite, which is a fibrous or acicular variety of hornblende.

Vein - A thin sheet-like intrusion into a fissure or crack, commonly bearing quartz /a small vein or cluster of veins.

Volcanic - Descriptive of rocks originating from volcanic activity.

Volcano-sedimentary - A mix of rocks formed by volcanic and sedimentary processes.

2. FORWARD-LOOKING/SAFE HARBOUR STATEMENT AND FAIR DISCLOSURE STATEMENT

This MD&A may contain certain forward looking statements concerning the future performance of the business of Houston Lake Mining Inc. (“HLM”) or the (“Company”), its operations and its financial performance and condition, as well as management’s objectives, strategies, beliefs and intentions. These forward-looking statements are based on information currently available to the Company and the Company provides no assurance that actual results will meet management’s expectations. Forward-looking statements include estimates and statements that describe the Company’s future plans, objectives or goals, its ability to access capital, the speculative nature of mineral exploration and development, fluctuating commodity prices, competitive risks and reliance



on key personnel, and include words to the effect that the Company or management expects a stated condition or result to occur. This list is not exhaustive of the factors that may affect any of the Company's forward-looking statements. Statements relating to estimates of reserves and resources are also forward-looking statements as they involve risks and assumptions, including but not limited to assumptions with respect to future commodity prices and production economics, that the reserves and resources described exist in the quantities and grades estimated and are capable of economic extraction. Forward-looking statements may be identified by such terms as "believes", "anticipates", "expects", "estimates", "may", "could", "would", "will", or "plan". All forward-looking information is inherently uncertain and subject to risks, uncertainties, and a variety of assumptions to address future events and conditions. These and other factors should be considered carefully and readers should not place undue reliance on the Company's forward-looking statements. The Company does not undertake to update any forward-looking statement that may be made from time to time by the Company or on its behalf, except in accordance with applicable securities laws.

3. HISTORICAL RESOURCE ESTIMATES

HLM's projects include properties with historical resource estimates which are not compliant with National Instrument 43-101 ("NI 43-101"). These estimates are sourced from various government and company archives which provide information on the geology and extent of the mineralization. A "qualified person" has not done sufficient work to classify the historical estimate as a current mineral resource or mineral reserve. Houston is not treating historical estimates as current mineral resources or mineral reserves as defined by NI 43-101 and historical estimates should not be relied upon.

4. INTRODUCTION

HLM is a "reporting issuer" in the Canadian provinces of Ontario, Alberta and British Columbia. The Company's common shares trade on the TSX Venture Exchange ("TSX.V") under the symbol HLM.

HLM is a specialized mining exploration company actively engaged in the acquisition, exploration and development of hard rock, high grade, rare metal resource properties. HLM maintains a particular focus on Lithium, Cesium, Tantalum and Rubidium by its world class expertise and experience exploring for LCT pegmatites.

Currently, HLM is actively exploring its 100% owned and optioned PAK Rare Metals Project in northwestern Ontario, Canada.

The Company's shares are listed on the TSX Venture Exchange under the symbol HLM and the head office is located in Val Caron, just north of Sudbury, Ontario.

The following management discussion and analysis should be read in conjunction with the Corporation's audited annual financial statements for the year ended March 31, 2013 and 2012 which comply with International Financial Reporting Standards ("IFRS"). All amounts are expressed in Canadian dollars unless otherwise stated.

Additional information regarding the Company can be found on SEDAR at www.sedar.com. The Corporation's web site is www.houstonlakemining.com.

5. ASSET OVERVIEW

Since start-up in 1995, Houston Lake acquired 100% interests in 9 properties that comprise the Company's three significant project groups: the West Cedartree Gold Project near Kenora, the Tib Lake PGM Project near Thunder Bay and the Pakeagama Rare Metals Project near Red Lake.



Figure I: Houston's Key Exploration Projects

2012 was a very difficult year for small and large cap companies. Although gold prices have remained somewhat stable after a short period of decline, most retail and institutional investors have remained on the sidelines. Share prices of many exploration and gold producing companies have fallen dramatically and in many instances, reached all-time lows. Needless to say, small caps have been hit the hardest. Many well-funded companies have had to suspend and defer exploration programs whereas others have been looking for strategic partners. Raising monies for exploration has become very dilutive and somewhat counter-productive insofar as share prices did not reflect the increase in value from having brought assets closer to the production stage. Overall, the market has paid little attention to the investment monies spent to enhance shareholder value.

HLM has taken decisive action on past interests expressed by North American Palladium Ltd. (TSX: PDL, NYSE Amex: PAL) in our Tib Lake PGM Project and Coventry Resources Ltd. (ASX:CVY, TSXV: CYY) in the West Cedartree Gold Project. HLM has been neighbours at our respective exploration projects with these companies for several years. Both PDL and Coventry put forward proposals to purchase our platinum group metals and gold assets and we have taken advantage of these interests and turn this opportunity into cash and to increase our focus on our PAK Rare Metals Project in northwestern Ontario.

6. PAK RARE METALS PROJECT

In January of 1999, the Ontario Geological Survey (OGS) published an article on a new discovery with potential for economic tantalum, cesium, rubidium and lithium mineralization at Pakeagama Lake, located about 160km north of Red Lake in northwestern Ontario ([Figure II](#)).



Figure II: Project Location Map

In late March of 1999, HLM entered into an option agreement to earn a 100% interest in the property. The area in the vicinity of Pakeagama Lake was mapped by Dr. Denver Stone of the OGS in the early 1990's. He identified a series of fertile granites and took a number of samples from the Pakeagama Lake pegmatite which contained highly anomalous lithium, tantalum and cesium values. Based on this initial field work, Dr. Fred Breaks of the OGS conducted three weeks of detailed mapping and sampling in 1998. Dr. Breaks identified a strongly zoned, complex petalite subtype pegmatite. The combination of size, accessibility and the presence of both geochemical indicators and rare metal ore minerals, provide the project with high potential for economic rare metal mineralization.

In 2011 HLM, under new management entered into an agreement with two private individuals to acquire 3 mining claims (each 16 unit claims) collectively called the Pakeagama South-East Property. The property optioned covers the strong potential of a south to east extension of the Pakeagama rare metal pegmatite. In late 2011, HLM commenced the field work for an Mobile Metal Ion (MMI) geochemical survey, including an initial orientation line over the Pakeagama Lake pegmatite on the PAK Lake, and PAK South-East Properties.

In 2012, HLM expanded the PAK Rare Metals Project to the north-west of the exposed pegmatite by staking. The company also completed analysis of the MMI geochemical survey sample collection conducted in 2011 on the project. In late 2012, HLM conducted a Phase I, 91 Channel Sample program in order to confirm historical results and to add additional channel samples while further sampling by distinct geologic zones. Under HLM's QA/QC procedures, sample blanks, along with tantalum, lithium, rubidium and cesium standard samples were routinely inserted into the sample stream.

In early 2013, HLM disseminated results from the Phase I Channel Sample program up to 4.74% Li₂O over 15 meters, 192 ppm Ta₂O₅ over 14 meters, and 0.53% Rb₂O over 14 meters 3 three distinct pegmatite zones.

6.1 Project Description and Location

The PAK Rare Metals Project is located 160 kilometres north of Red Lake, Ontario in the Red Lake Mining Division. The Project is

composed of 224 units for a total of 3,584 hectares (8,856 acres) as can be seen in [Figure III](#). The Pakeagama pegmatite is located on the Pakeagama Lake Property and is owned 100% by the company, and it is subject to a 2.5% net smelter return, with HLM maintaining the option to purchase partial royalties.




Figure III: PAK Rare Metals Project Claim Locations

An elongate, approximately 2.5 by 15 kilometre, Pakeagama Lake peraluminous granite and mica pluton that trends northwest-southeast was emplaced immediately adjacent to and parallel to the Sachigo-Berns River Sub-provincial boundary of the Superior Province of the Canadian Shield. It was emplaced along the unconformable contact between metasedimentary rocks and metavolcanic/metasedimentary rocks of Achaean age. The north-western portion of the granite is present in the south-eastern quadrant of claim KRL1232441 (Pakeagama Lake Property). The granite is defined as fertile granite; the granite is a potential source of the rare earth elements found in the Pakeagama Lake Pegmatite.

The economically important Pakeagama pegmatite is located within the boundaries of claim KRL 1232441 at the northwest end of the Pakeagama Lake Granite. The pegmatite is a highly evolved, complex-type, petalite subtype pegmatite with highly anomalous to economically significant values of tantalum, cesium, rubidium and lithium (F.Breaks, A.Tindle, and S.Smith, 1999a). The pegmatite is divided into three distinct zones: the upper intermediate zone (UIZ), the central intermediate zone (CIZ), and the lower intermediate zone (LIZ). Metasedimentary rocks in unconformable contact with metavolcanic/metasedimentary rocks underlie all other parts of the claim KRL1232441.

6.2 Accessibility, Climate, Local Resources, Infrastructure and Physiography

The project is located in the Pakeagama Lake township, Red Lake Mining district of north-western Ontario. The National Topographic System map sheet reference is 53C/11, and the Pakeagama Pegmatite is approximately centred at 52°36'N latitude and 93°23'W longitude ([Figure IV](#)).




Figure IV: Project Regional Geology

One hundred and sixty kilometer float plane access is available from Red Lake to the west side of Pakeagama Lake where there is a summer camp site on a point of land. Red Lake is located one hundred and seventy kilometres north of Vermillion Bay on Highway 105. Vermillion Bay is situated on the Trans Canada Highway (Hwy. 17) between the centres of Dryden and Kenora. Pakeagama Lake is located in a relatively isolated area of north-western Ontario. A winter road passes adjacent to the PAK Rare Metals Project claim group, whereby a winter camp site is located. Bear Skin Airways services the nearby communities of Deer Lake, North Spirit Lake, and Sandy Lake.

The Pakeagama Lake area lies at the northern boundary of the Lac Seul Upland ecoregion and the southern boundary of the Hayes River Upland ecoregion of the Boreal Shield. The region is classified as having a subhumid mid-boreal ecoclimate (Ecological Stratification Working Group, 1998). The Pakeagama Lake area lies between four data stations which are located at Island Lake (Manitoba), Red Lake (Ontario), Pickle Lake (Ontario), and Big Trout Lake (Ontario) for which extensive climatic records are available from Environment Canada. The average mean annual temperature is -0.9°C . The average daily temperature in summer is from 8.9°C to 19.2°C while in winter the average range is from -20.3°C to -11.6°C . Average yearly precipitation for the area is 655 mm. The dominant vegetation is coniferous forest. The area has seen a forest fire in approximately 1995 and deadfall inhibits easy foot travel. Another forest fire burned a portion of the Pakeagama Lake claim in 2008, including the area around the Pakeagama pegmatite. Higher elevations are covered by stands of jack pine up to 20 feet high while swampy areas are dominated by black spruce. Upland areas are covered with discontinuous deposits acid, sandy tills while thin lacustrine clay deposits may cap the tills in low-lying areas. Wildlife includes wolf, lynx, moose, black bear, red squirrel, snowshoe hare, beaver, mink, fisher, marten, and ermine. The area lies north of the Government of Ontario's "Lands for Life" and "Living Legacy" programs of conservation and wilderness preservation. The claim contains a series of ridges with the extensive development of cliff faces that parallel the general regional strike of the geology. The ridges are separated by relatively flat, muskeg terrain. Infrastructure is absent except for the winter road, which is located adjacent to the claim group.



6.3 History

The first geological reconnaissance mapping of the region was undertaken by A. P. Low of the Geological Survey of Canada in 1886. Additional geological surveys were carried out by G. V. Douglas (1925) and M. E. Hurst (1928) of the Ontario Department of Mines. Most of the exploration activity in the region centred on the Favourable and Setting Net Lakes area located 25 to 40 km to the northwest of the property. Prospecting by K. C. Murray in 1927 identified gold-bearing mineralization in the Favourable Lake area. The gold property was developed as the Berens River Mine and produced 4,451 kg Au, 160,926 kg Ag, 2,770 t Pb and 815,147 kg Zn from 508,665 tons of ore between 1939 and 1948 (Stone, 1998). Subsequent exploration by Golsil Mines Limited, Zahavy Mines Limited, Getty Mines Limited and Noramco Mines Ltd. was carried out until the early 1990's. The "Wynne" gold showing is located on the northern shore of Pakeagama Lake. However, there are no records of any assessment work having been filed from the vicinity of Pakeagama Lake. The Pakeagama Lake area was covered by an airborne reconnaissance gamma-ray spectrometer survey in 1977 as part of a regional coverage program (OGS-GSC, 1979). The survey was flown at a 400-ft elevation with 5-km line spacing and a 2.2 km station interval. No significant radiometric anomalies were detected in the immediate vicinity of Pakeagama Lake. Geological mapping of portions of the region was carried out by Ayres (1970, 1972a). He noted spodumene in a pegmatite dike and holmquistite within granitic rocks near Setting Net Lake (25 km WNW of Pakeagama Lake). A grab sample from the pegmatite dike returned 0.52% Li (Ayres, 1972b). Recent mapping of the region was carried out by D. Stone of the Ontario Geological Survey in 1990 (Stone et al, 1993, Stone, 1998). Tourmaline-rich samples taken from the vicinity of Pakeagama Lake returned anomalous Li, Cs, Ta and Be during this work. Detailed follow-up work was carried out by Breaks et al (1999a) in the vicinity of Pakeagama Lake. Five rare metal mineral occurrences were detected over a 35 km segment of the Bear Head Fault Zone. However, the Pakeagama Lake pegmatite occurrence became the predominant focus of detailed work. G. Anthony mapped the pegmatite and adjacent rocks during 1999. Check assays of samples collected by the Ontario Geological Survey were also completed. HLM completed a program of line cutting, ground geophysics, prospecting, rock saw channel sampling and overburden removal on claim 1232441 (16 claim units), between May 12 and July 25, 2001 (not NI 43-101 compliant). A study performed by Dr. R. Ken Germundson, P. Geo determined that the tantalum oxide values for samples collected for HLM were about two thirds as great as the Ontario Geological Survey samples. There was a close agreement between the three sets of data for the average content of cesium, although no rare metal standards were inserted into any of the government, or HLM sample stream.

6.4 Geological Setting

i. Regional Geology

The Pakeagama pegmatite, which contains rare metals, is located in HLM's claim 1232441 (Pakeagama Lake Property). Bands of iron formation are present within both of the sequences of metamorphic rocks that are present in the project. As can be seen in [Figure IV](#), the project is located adjacent to the boundary between the Berens River and Sachigo Subprovinces of the Archean-age Superior Province of the Canadian Shield. These subprovinces comprise a series of relatively isolated volcano-sedimentary (greenstone) belts surrounded by extensive granitic and gneissic suites of rock. They are separated by the Bearhead Lake Fault Zone.

Two of the greenstone belts that are located along the contact between the two sub provinces are: the Favourable Lake greenstone belt located towards the northwest, and the North Spirit Lake greenstone belt located towards the southeast of the property. The belts are connected through the Pakeagama Lake area by the Bearhead Fault. The main assemblages of volcanic and sedimentary rocks that are identified in each belt are, in part, correlated between the two belts. The assemblages of the Favourable Lake and North Spirit Lake areas are predominantly in greenschist facies of metamorphism. However, a transition to amphibolite facies in the greenstone belts occurs as the Bear Head Fault Zone is approached. Amphibolite facies is the predominant metamorphic grade in the Pakeagama Lake vicinity.

The Bear Head Fault is the dominant structural feature in the region and has been traced for over 140 km from NW-SE. The fault is composed of a several hundred metre thick zone of mylonite. The presence of cataclastites, tension gashes unfilled by vuggy quartz-epidote-adularia and potassic alteration indicate that brittle deformation has been superimposed on the mylonites. A dextral transcurrent dislocation of the Bear Head Fault has been interpreted from microstructures. The regional gneissosity trends NW-SE and generally are steeply dipping inward towards the core of the volcano-sedimentary assemblage in the vicinity of Pakeagama Lake. The Bear Head Fault Zone is the locus for a peraluminous suite of granitic plutons. Five major plutons of the two-mica granites (fertile granites) are documented over the 140-kilometre strike length of the fault. Fertile granites are interpreted to be the parental rocks that give rise to rare metal pegmatites.

ii. Property Geology



The exposed pegmatite on the project has been classified into three main lithological domains. The northeast third of the Pakeagama Lake claim block is dominantly of metasedimentary origin and composed of pelitic sediments, iron formation and conglomerate. The southwest third is comprised dominantly of mafic metavolcanic and related metasedimentary rocks. The Pakeagama Lake pluton dominates the southeastern portion and appears to pinch out in the south-central part of the property. The geology of the area of claim KRL 1232441 is presented in Map P. 3224 (Whiteloon Lake) of the Ontario Geological Survey (Stone and Fitzsimon, 1993). The property is underlain by the north-western extension of the North Spirit Lake greenstone belt. The greenstone rocks, which are approximately 2 kilometres wide in the vicinity of the property, are bounded to the north by biotitic tonalities and granites of the Whiteloon Lake Batholith (Sachigo Subprovince of the Superior Province of the Canadian shield) and to the south by gneissic granodiorites and granites of the Bearhead Lake Batholith (Berens River Subprovince of the Superior Province).

iii. Pakeagama Lake Pegmatite Geology

Granitic pegmatites are those pegmatites derived from granitic intrusions. They may be divided into classes, types and families on a combination of criteria. The criteria include minor elements present and the relative concentrations of minor elements (i.e. tantalum to niobium), geological setting and metamorphic grade. The rare element class of granitic pegmatites is the only grouping that has significant economic potential. A zoned pegmatite body occurs near the north western margins of the Pakeagama Lake pluton in the south central portion of the property. Muscovite and tourmaline-bearing pegmatites and aplites may occur up to 1 km from the main pegmatite mass (F. Breaks, A. Tindle, and S. Smith, 1999a). The Pakeagama Lake pegmatite is the second largest complex-type petalite sub-type pegmatite dike in Ontario (F. Breaks, A. Tindle, and S. Smith, 1999a). The relatively fresh-appearing pegmatite dike has irregular steeply dipping contacts with the weakly foliated garnet-muscovite-biotite granite host rock. A 130° strike is inferred from the coincidence of the exposed 260 metre strike length, the weak foliation in the host granite and the general trend of the Bear Head Fault. The pegmatite is open along strike in both directions.

A zoned pegmatite body occurs near the margins of the Pakeagama Lake pluton in the south central portion of the property. The pegmatite that is exposed at surface is the only portion of the project that has been mapped and recently sampled in detail. The current known size of the rare-metal pegmatite is largely within the confines of the Pakeagama Lake granitic pluton. On surface, a banded iron formation forms an apparent northern boundary to the pegmatite. Three pegmatite zones have been depicted and confirmed by HLM on the known exposure. From northwest to southwest and perpendicular to the strike of the pegmatite the zones include a Central Intermediate Zone (CIZ – tantalum and rubidium enriched), Upper Intermediate Zone (UIZ - lithium and rubidium enriched), and the Lower Intermediate Zone (LIZ – lithium and rubidium enriched).

Central Intermediate Zone (CIZ)

The Central Intermediate Zone (CIZ) is located in the upper portions of the pegmatite. This altered zone, as sampled by Channel 1, contains the highest tantalum grades found to date in the exposed pegmatite. As can be seen in [Figure IV](#) it is in contact with both the Upper Intermediate Zone and Upper Wall Zone, and persists to the southeast edge of the outcrop where it is believed the pegmatite continues under the till cover. Channel 19 is located in the central portion of the exposed CIZ where the zone consists of predominantly grey K-feldspar with minor lithian mica + quartz alteration. To the southeast the alteration process whereby the K-feldspar is altered to a mica+quartz mixture was sampled by Channel 7R. The samples from this channel consisted of similarly sized fragments of randomly oriented K-feldspar with varying degrees of mica+quartz alteration. In the adjoining area to the northeast of Channel 7R, the alteration process whereby the K-feldspars are more or less completely replaced with a lithian mica + quartz mixture is visible. In this area veinlets and patches of lepidolite occur.

Upper Intermediate Zone (UIZ)

The UIZ represents the lithium zone within the pegmatite and is dominated by "SQUI", a term used to describe an oriented spodumene + quartz intergrowth after primary petalite, with lesser grey K-feldspar and primary white spodumene in quartz. Phosphate minerals and lithian mica are the common accessory minerals. Channels 3A, 3B and 4 with a composite grade of 4.74% Li₂O represent a 15 metre section across the UIZ in the southeastern part of the exposed pegmatite. Channel 18 is located approximately 55 metres northwest along strike from Channels 3A/3B/4.

Lower Intermediate Zone (LIZ)

The LIZ comprises a good portion of the exposed pegmatite and is considered an intermediate stage zone historically carrying significant lithium, tantalum and rubidium. The zone is comprised of predominantly K-feldspar, Na-feldspar, quartz, SQUI and



lithian muscovite. The zone has undergone both plastic and brittle deformation. Taken together the three channels represent a 23 meter cross-section of the LIZ with a composite grade of 2.27% Li₂O. When considering the LIZ and the high grade UIZ as outlined in Table 1, the total distance between the two sets of channels is approximately 190m where the pegmatite zones remain open to both the southeast and northwest.

6.5 Geological Target: Rare Metal Target (Tanco Mine)

The word 'pegmatite' stems from Homeric Greek meaning 'to make stout or rigid by binding together'. The crystallization of fertile granites yields a fractionated residual magma which is enriched in lithium, rubidium, cesium, thallium, tantalum, hafnium, gallium, germanium, boron, fluorine and phosphorus relative to that contained in the fertile granite. This fractionated residual magma is expelled out from the fertile granite source into the surrounding metasedimentary and metavolcanic rocks where it cools and consolidates to form a group of pegmatite bodies.

The pegmatite group derived from the fertile granite parent shows a progressive fractionation trend and chemical changes with distance from the fertile granite source. "There have been many models invoked on the study of rare-element pegmatites, but regardless of which is used the pegmatite-forming process is dynamic and highly process-dependent, involving a rather delicate thermal and chemical balance" (Jahns 1953) to explain the many complex textural and mineralogical features of pegmatites. The most fractionated batches of pegmatite are the most mobile (enriched in boron, water, fluorine, phosphorus, lithium, rubidium, cesium, tantalum and beryllium) and tend to migrate the furthest from the fertile granite parent. The most fractionated of these pegmatites are the complex type pegmatites which include the spodumene, petalite, lepidolite, and amblygonite subtypes. Individual pegmatites are generally small and range from several metres long and 1 metre thick to economically interesting bodies hundreds of metres long and tens of metres thick.

The internal structure of pegmatites varies from homogenous to zoned. Zoning or the lack of it provides the starting point for internal anatomy. Zoning is largely what distinguishes pegmatites from other ordinary plutonic igneous rocks. It is manifested by variations in the spatial distribution of grain size, mineral assemblage, crystal habit, or rock fabric. Zoned pegmatites may have up to nine different units with variable textures and mineral modes (Cerny, 1991b, Cerny et al, 1996). The most evolved zonation patterns are encountered in highly fractionated intrusions with complex rare metal mineralization (i.e. Tanco).

The most striking geochemical anomalies of the Tanco body are its high Cesium and Phosphorus contents, the hallmark of a pelitic metasedimentary source, and high tantalum.

6.6 Exploration Programs

i. Past Exploration

In 1998 and 1999 Dr. Fred Breaks of the Ontario Geological Survey and Dr. Andy Tindle studied the Pakeagama Pegmatite. Approximately 2186 analyses have been conducted to establish that the Pakeagama Rare Metals Pegmatite is potentially a world-class pegmatite suggesting the presence of a Tanco type mineralizing system. The Pakeagama Lake pegmatite has a width which varies from 30 to 125 metres and a strike length of at least 260 metres (open in both directions) which may extend another 300 metres to a historical 700 g/t tantalum oxide occurrence in an aplite dyke. In Summary of Field Work 1999, OGS Open File Report 6000 dated November, 1999, Dr. Breaks et al conclude: "The detailed documentation of a variety of tantalum-rich minerals coupled with the presence of pollucite (main cesium ore mineral) renders the Pakeagama Lake pegmatite and adjoining area one of the best exploration targets for tantalum and cesium in Northwestern Ontario".

ii. Recent Exploration

In June 2011, HLM conducted an MMI orientation soil survey over the exposed Pakeagama Lake pegmatite. The results of the survey identified 15 elements that had a definitive response to the pegmatite, and specifically established a characteristic cesium, lithium, and rubidium soil MMI response. It also allowed the Company to define the best soil sample depth range for future MMI surveys on the property. Later in 2011, the Company performed a detailed reconnaissance MMI soil survey of the Pakeagama Lake, and PAK South-East claims. The regional survey returned very encouraging MMI results. Specifically, elevated cesium, lithium and rubidium MMI values are concentrated both to the northwest and southeast directions within the property and coincident with the assumed orientation of the Pakeagama Lake pegmatite. The MMI geochemical signatures are of similar amplitudes to that of the orientation survey over the Pakeagama Lake pegmatite itself. The objective of the MMI geochemical program is to test a cost effective geochemical survey tool for further definition of the Pakeagama Lake rare metal pegmatite, or

other similar pegmatites in central Canada.

In September 2012, HLM conducted a Channel Sample Program on the Pakeagama Lake pegmatite. The 91 sample program was conducted in order to confirm historical results and to add additional samples while further sampling by distinct geologic zones (UIZ, LIZ, and CIZ). The program identified high grade lithium, tantalum, and rubidium up to 4.74% Li₂O over 15 metres in three distinct pegmatite zones, and 14 metres of 192ppm Ta₂O₅, including 270ppm over 6 metres, and 0.53% Rb₂O in one of these zones (Figure V).




Figure V: Channel Samples & Detailed Geology

On March 1, 2013, HLM announced the commencement of a Phase I, 6-hole diamond drill program as a follow-up and focus on drill targets derived from the high grade mineralization encountered in the UIZ, LIZ, and CIZ zones sample with the 2012 Channel Sample Program. The objectives of the program are (1) to determine the orientation, thickness and zonation of the pegmatite, and (2) to refine the mineralogical characterization of these zones by better establishing their lithium, tantalum, cesium and rubidium potential. The dissemination of results are expected to commence in the spring of 2013.

7. WEST CEDARTREE GOLD PROJECT (WCGP)

7.1 Sale of WCGP

On January 15, 2013 HLM announced the signing of a definitive agreement with Coventry Resources Limited (TSX.V: CYY, ASX: CYY) ("Coventry") who owns the Cameron Gold deposit, located 12km away from the West Cedartree Gold Project, as well as their Rainy River Project located 80km away. Under terms of the agreement, HLM received total consideration at the time which was valued at approximately \$1.22 million, allocated as follows (all currency amounts in Canadian dollars, unless otherwise indicated):

- \$100,000 in cash, on execution of the binding Heads of Agreement;
- \$400,000 in cash, on execution of a Definitive Sale and Purchase Agreement (the "Agreement"),



- 1.935 million Coventry shares, representing 2.8% of the outstanding shares of Coventry, (on execution of the Agreement) with a value of approximately \$716,000. Coventry shares closed on the TSX.V at CAD\$.37 on January 11, 2013); and,
- A 2.5% net smelter return (“NSR”) royalty on the West Cedartree Claim Block, representing 20% of the West Cedartree Gold Project’s total area, including the newly discovered Robertson Prospect. The NSR does not have a buy-back clause, however should HLM seek to sell the royalty, Coventry will maintain the first right of refusal.

HLM is very pleased to have taken a significant step with such an aggressive, capable and well respected exploration company as Coventry and believe that they have the desire and expertise to advance their project to the benefit of both companies, and the West Cedartree Gold Project. Coventry’s significant Cameron Gold deposit, coupled with their strong financing capabilities provide the opportunity for HLM shareholders to capture value with cash and liquid assets in order to focus on our PAK Rare Metals Project in Northwestern Ontario; to retire some outstanding debt; and to continue to participate in the upside of their Cameron Gold and Rainy River Projects.

8. TIB LAKE PGE PROJECT

8.1 Definitive agreement on option to purchase Tib Lake PGE Project with North American Palladium

A definitive agreement was reached with North American Palladium Ltd. (TSX: PDL, NYSE Amex: PAL) for the option to purchase the Tib Lake PGE Property on May 8, 2013. The agreement provides NAP with an option to earn a 100% interest in the 2,464-hectare Tib Lake PGE property from HLM after making staged payments totaling CAD\$450,000 to HLM and incurring exploration expenditures of CAD\$1.6 million over five years. Upon completion of the earn-in, the property will have a residual 2.5% net smelter return royalty (NSR) in favour of a third party or HLM (depending on the claim). HLM will also retain the right to buy back 1% of the third party NSR for CAD\$1 million and to purchase the remaining 1.5% of the third party NSR on such terms and conditions that the two parties may agree on. On claim areas where HLM is a royalty holder, NAP will have the right to purchase 1% of the NSR for CAD\$1 million.

8.2 Tib Lake PGM Property Details

The 2,496 hectare Tib Lake PGM property (2.5% NSR) is located just 15km from North America Palladium’s 15,000 tonne per day mill at Lac des Iles Mine (“LDI”). The property covers 100% of the highly prospective Lower Zone of the Tib gabbro, the largest exploration target for platinum group metals (“PGM’s”) in the vicinity of the LDI mine.




Figure VI: Basic Geology of the Tib Lake and Lac Des Iles Intrusions

Four PGM occurrences have been documented in the Lower Zone: the Jewelweed, the Road, the West Shore and the Kuhner. The Kuhner Occurrence is the main site of previous exploration. Previous drilling intersected core lengths of: 18.5m grading 1.58 g/mt PGM's (Pt-Pd-Au), 22.0m of 1.46 g/mt PGM's and 14.5m grading 1.12 g/mt PGM's.

The exploration potential of the Tib Lake PGM Project evolves from a synthesis formed from the generalized PGM ore deposit model, comparative observations of the geology of the Lac Des Iles mine vicinity and the Tib Lake gabbro, and previous exploration conducted on HLM's Tib Lake PGM project. Over the past Nine Months NAP has increased its land position in the LDI region through the acquisition and option agreements of several high-potential PGE properties, all of which are located less than 30 kilometres from the LDI mill and whereby Tib Lake sits at the top of that list through both its size and proximity. NAP believes that Tib Lake relates to the same magmatic event that produced the LDI complex and its world-class palladium resources. HLM believes that the knowledge gained by NAP from two decades of exploration and mining at LDI will be applied to the Tib Lake intrusion, and based on their insatiable requirement of feed for their underutilized 15,000 tonne-per-day mill HLM is poised for potential upside. NAP has recently released "An aggressive surface exploration program involving trenching, mapping, prospecting and geochemical surveys was initiated at the beginning of July. The discovery of new PGE resources stemming from a sustained investment into PGE greenfields exploration remains a strategic priority for NAP."

9. EXPLORATION STANDARDS

HLM conducts exploration activities in accordance with "Exploration Best Practices Guidelines" established by the Canadian Institute of Mining, Metallurgy and Petroleum (CIM) and conforms to NI 43-101 standards. Houston Lake's exploration programs are managed by Qualified Persons as defined by NI 43-101.

10. EXPLORATION ACTIVITIES AFFECTING THE BALANCE SHEET

10.1 Results of Operations for the three-month period ended March 31, 2013

The Company's loss for the three-month period ended March 31, 2013 (the "Current Period") was \$7,219,399 or \$.12 per share as compared with a loss of \$345,895 or \$.008 per share for the three-month period ended March 31, 2012 (the "Comparative Period").



The significant increase in losses is attributed primarily to the loss on the sale of the West Cedartree Gold Project.

General and administrative expenses were \$21,448 lower in the Current Period at \$13,849 compared with \$35,297 in the Comparative Period. Costs were reduced as a result of decreased exploration expenditures (\$271,478 versus \$378,015) due to lower exploration spending on the Company's properties. The Company had decreases in professional and consulting fees (\$37,634 versus \$69,134) also due to lower exploration expenditures by the Company.

The Company also realized a loss on sale of exploration and evaluation asset during the last quarter of \$7,531,341 related to the sale of the West Cedartree Gold Project to Coventry Resources Ltd.

10.2 Results of operations for the year ended March 31, 2013

During the year ending March 31, 2013 HLM invested \$331,243 in the mining properties. \$24,000 of that amount was spent on the PAK Rare Metals Project as a part of the acquisition costs associated with the option on the PAK Southeast Property and the balance of \$307,243 was spent on the 2012 Channel Sample Program and part of the Phase I, 6-hole diamond drill program.

For the year ended March 31, 2013 HLM incurred a net loss of \$7,589,439 (\$0.13/share) compared to a net loss of \$846,315 (\$0.02/share) for the same period ending in 2012. The significant net loss for 2013 is primarily due to the loss on the disposal of mining properties encompassed by the West Cedartree Gold Project sale to Coventry Resources Ltd.

11. FINANCING ACTIVITIES

Ongoing exploration was funded in part by the divestiture activities by the company with the West Cedartree Gold Project, Tib Lake PGM Project and \$200,000 financing in August 2012. Furthermore, the re-structuring of HLM has enabled it raise \$431,000 with the issuance of 8.9 million common shares and warrants in order to further exploration on the PAK Rare Metals Asset. The \$433,000 financing was a subsequent event as it was finalized on April 09, 2013.

12. CAPITAL STOCK

HLM had 63,019,056 shares outstanding at March 31, 2013. 8,900,000 shares were issued during the year ended March 31, 2013. No options were exercised during the twelve month period.

13. LIQUIDITY AND WORKING CAPITAL

Current assets of the Company were \$550,028 as at March 31, 2013. As a result of the divestiture activities of HLM, an increase from March 31, 2012 of \$517,917 has been realized.

Net working capital totaled \$-207,510 as at March 31, 2013 compared to \$-705,843 at the March 31, 2012 year end due primarily from the cash generation due to the sale of the West Cedartree Gold Project. Current liabilities increased slightly from \$737,954 at March 31, 2012 to \$757,538 for the current period. Of this amount, \$496,614 is due to corporations controlled by a director of HLM.

The Company's current rate of cash consumption, excluding expenditures on work programs or cost of financings is approximately \$30,000 per month. On average the Company's rate of cash consumption for property expenditures (PAK Rare Metals Project) was \$28,000.

All of the Company's properties are 100% owned with minimal holding costs except the 100% optioned PAK Southeast Property. A total of \$24,000 in option payments were made in the fiscal year of 2013 on the PAK Southeast Property.

Future cash generation will be supplied by the account receivable of \$310,000 from North American Palladium Ltd. over the next three years, as well as the raising of monies through the capital markets.

14. OFF-BALANCE SHEET ARRANGEMENTS

As at March 31, 2013 the Company does not have off-balance sheet arrangements.

15. ASSET-BACKED COMMERCIAL PAPER

As at March 31, 2013 the Company does not have and has never had any exposure to asset-backed commercial paper.

16. RELATED PARTY TRANSACTIONS

During the year ended March 31, 2013, the Company incurred a total of \$10,000 in cash expenditures with a company controlled by an officer.



Included in accounts payable is \$327,785 (2012 of \$168,112) owing to companies controlled by a director, \$2,191 (2012 - \$62,415) owing to corporate directors and \$53,906 owing to corporate officers for consulting fees.

A loan payable of \$114,923 (2012 - \$388,648) is owed to a corporation controlled by a director and bears interest at the Royal Bank prime lending rate plus 2%, is unsecured and is due on demand. The loan payable costs are primarily related to the carrying costs of the director controlled company for consulting wages related to an officer of HLM. Interest in the amount of \$4,452 (2011 - \$4,452) has been accrued and remains unpaid at year –end.

17. OUTLOOK

HLM will continue to focus on exploration and development objectives on the PAK Rare Metals Project.

With the TSX Venture Index experiencing negative returns year to date, the lengthy Eurozone debt crisis, signs of Southeast Asian growth slowing, and dropping commodity prices, the raising of capital is proving difficult for exploration companies. Management believes that such an environment instigates asset and/or corporate consolidation, especially for small cap companies. HLM's divestiture actions are within response of current corporate debt, capital market stresses and the above mentioned industry trend of consolidation. Streamlining the asset base through divestiture has reduced working capital requirements, and also providing a re-branding of the company in the rare-metal category of exploration.

18. RISKS AND UNCERTAINTIES

HLM's may be exposed to risks of varying degrees of significance which could affect its ability to achieve its strategic objectives. The main objective of the Company's risk management processes is to ensure that the risks are properly identified and that the capital base is adequate in relation to those risks. Risks include metal price fluctuations and the low success rate for the discovery of new deposits. Industry competition and lack of funding may also limit opportunities. Future political, regulatory and environmental changes could affect any aspect of the Company's business including property title, taxation, aboriginal issues and environmental protection. More detail of the principal risks to which the Company is exposed to are described below:

i. Credit Risk

Credit risk is the risk of loss associated with a counter party's inability to fulfil its payment obligations. The Company's credit risk is primarily attributable to cash and cash equivalents and accounts receivable. Cash and cash equivalents consists of cash on hand deposited with reputable financial institutions which is closely monitored by management. Accounts receivable includes HST receivable and is subject to CRA's assessment prior to receipt. Management believes credit risk with respect to cash and cash equivalents and accounts receivable is low.

ii. Liquidity Risk

The company has no history of profitable operations and its present business is at an early stage. As such, the Company is subject to many risks common to such enterprises, including under-capitalization, cash shortages and limitations with respect to personnel, financial and other resources and the lack of revenues. There is no assurance that the Company will be successful in achieving a return on shareholders' investment and the likelihood of success must be considered in light of its early stage of operations.

The Company has no source of operating cash flow and no assurance that additional funding will be available to it for further exploration and development of its projects when required. Although the Company has been successful in the past in obtaining financing through the sale of equity securities or joint ventures, there can be no assurance that the Company will be able to obtain adequate financing in the future or that the terms of such financing will be favorable. Failure to obtain such additional financing could result in the delay or indefinite postponement of further exploration and development of its properties. However, the Company ensures that there is sufficient cash and other short-term assets readily convertible into cash in order to meet its liabilities when they come due. The Company's cash is held in business accounts with a Canadian bank. Management believes that liquidity risk is low.

iii. Asset Exploration Risk

The Company's Property interests are located in remote, undeveloped areas and the availability of infrastructure such as surface access, skilled labour, fuel and power at an economic cost, cannot be assured. These are integral requirements for exploration, development and production facilities on mineral properties. Power may need to be generated on site.

Resource acquisition, exploration, development, and operation, is a highly speculative business that involves significant risks, which even a combination of careful evaluation, experience and knowledge may not eliminate. While the discovery of precious metals and other minerals may result in substantial rewards, few properties that are explored are ultimately developed into producing mines. Major expenses may be required to locate and establish economically viable mineral deposits, to develop



metallurgical processes and to construct mining and processing facilities at a particular site. It is impossible to ensure that the acquisition, exploration or development programs planned by the Company will result in a profitable commercial mining operation. The potential for any project to eventually become an economically viable operation depends on numerous factors including: the quantity and quality of the minerals discovered if any, the proximity to infrastructure, metal and mineral prices (which vary considerably over time) and government regulations. The exact effect these factors can have on any given exploration property cannot accurately be predicted but the effect can be materially adverse.

The mineral industry is intensely competitive in all its phases. The Company competes with many other mineral exploration companies who have greater financial resources and technical capacity.

iv. Commodity Price Risk

The market price of precious metals and other minerals is volatile and cannot be controlled. The current and expected future spot prices have a significant impact on the market sentiment for investment in exploration companies and may impact the Company's ability to raise equity financing for its ongoing working capital requirements.

v. Interest Rate Risk

The Company's cash is held in business accounts with nominal interest rates. Management considers interest rate risk to be low. The Company's loans bear interest at a variable interest rate. Interest on these loans could change due to changes in the market interest rate.

vi. Currency Risk

Currency risk is the risk that the value of a financial instrument will fluctuate due to changes in foreign exchange rates. The Company retains a US Bank Account with a nominal balance. Management considers currency risk to be low.

vii. Sensitivity Analysis

Based on management's knowledge and experience of the financial markets, the Company believes that a 10% movement in interest rates and foreign exchange rates that may reasonably be expected to occur over the next twelve month period will not have a significant impact on the Company.

viii. Management Risk

The Company is very dependent upon the personal efforts and commitment of its existing management. To the extent that managements' services would be unavailable for any reason, a disruption to the operations of the Company could result, and other persons would be required to manage and operate the Company.

The Company's directors and officers serve as directors or officers, or may be associated with other reporting companies or have significant shareholdings in other public companies. To the extent that such other companies may participate in business or asset acquisitions, dispositions, or ventures in which the Company may participate, the directors and officers of the Company may have a conflict of interest in negotiating and concluding terms respecting the transaction.

19. ADDITIONAL DISCLOSURE FOR VENTURE ISSUERS WITHOUT SIGNIFICANT REVENUE

Additional disclosure concerning HLM's general and administrative expenses and mineral property costs is provided in the Company's audited statement of loss contained in its audited financial statements for the year ended March 31, 2013.

20. EVALUATION OF DISCLOSURE CONTROLS AND PROCEDURES

Management is responsible for establishing and maintaining a system of controls and procedures over the public disclosure of financial and non-financial information regarding the Company. Such controls and procedures are designed to provide reasonable assurance that all relevant information is gathered and reported on a timely basis to senior management, including the President, Chief Executive Officer (CEO) and acting Chief Financial Officer (CFO) so that appropriate decisions can be made regarding public disclosure.

The system of disclosure controls and procedures includes, but is not limited to, the Company's Disclosure Policy and Code of Business Ethics, the effective functioning of Disclosure and Audit Committees, procedures in place to systematically identify matters warranting consideration of disclosure by the Disclosure Committee and verification processes for individual financial and non-financial metrics and information contained in annual and interim filings, including financial statements, MD&A filings and other documents and external communications.



As required by CSA Multilateral Instrument 52-109, Certification of Disclosure in Issuer's Annual and Interim Filings, an evaluation of the effectiveness of the design and operation of our disclosure controls and procedures was conducted, under supervision of Management, including the President, CEO and acting CFO. The evaluation included documentation review, enquiries and other procedures considered by management to be appropriate in the circumstances.

Based on that evaluation, the President, CEO and acting CFO have concluded that the design and operation of the system of disclosure controls and procedures was effective for twelve months ended March 31, 2013. The President, CEO and acting CFO are also required, under Multilateral Instrument 52-109, to file certifications of the annual filings. Copies of these certifications may be found on SEDAR at www.sedar.com.

HOUSTON LAKE MINING INC.

Henry J. Kloepper
CEO

July 24, 2013